

DATA LOOK JANUARY 2015 WISCONSIN TOBACCO PREVENTION & CONTROL

ALL TIME LOW RATES




2014 WI Youth Tobacco Survey



2013 Behavioral Risk Factor Surveillance System

POINTS OF CONCERN



14% of Wisconsin women smoke during pregnancy. 2012 Wisconsin Births and Infant Deaths

of high school students are exposed to secondhand smoke in their homes.

2014 WI Youth Tobacco Survey 15% of Wisconsin deaths are due to smoking. 2015 Burden of Tobacco in Wisconsin

TRENDS IN SMOKELESS & E-CIGARETTE USE

SMOKELESS

9.7%

of high school students currently use chewing tobacco, snuff, or dip. A 67% increase since 2012. (2012 rate: 5.8%)

2014 WI Youth Tobacco Survey

ELECTRONIC CIGARETTES

7.9%

of high school students have used e-cigarettes. 75% higher than the national average. (National average: 3.4%)

2014 WI Youth Tobacco Survey

4.7%

of Midwest adults have used e-cigarettes. (WI specific data not currently available)

National Adult Tobacco Survey, United States, 2012–2013

THE COST OF TOBACCO



3 billion

in health care costs are paid in Wisconsin as a result of diseases caused by smoking.

1.6 billion

in lost productivity costs because of illness and premature death from smoking-related illnesses.

2015 Burden of Tobacco in Wisconsin

\$528

estimated yearly health care cost of cigarette smoking for every man, woman, and child.

2015 Burden of Tobacco in Wisconsin

